


1. A method for the treatment or inhibition of hypercholesterolemia or hypertriglyceridemia, comprising the step of:
administering a composition comprising a therapeutically-effective amount of a compound of either Formula I:


a) Formula I



wherein R and R₁ are independently selected from the group consisting of H, lower alkoxy (C₁₋₆), or lower alkyl (C₁₋₆); R₂ is selected from the group consisting of aryl (C₆₋₁₂) and substituted aryl; M is hydrogen, sodium, potassium, ammonium, diethanolamine, cyclohexylamine, a naturally-occurring amino acid of MW less than 500 kD, lower alkyl (C₁₋₆), cycloalkyl, or aryl (C₆₋₁₂); and n is 0-5;

b) Formula III:

Formula III




wherein R and R₁ are independently selected from the group consisting of H, lower alkoxy (C₁₋₆), or lower alkyl (C₁₋₆); R₂ is selected from the group consisting of aryl (C₆₋₁₂) and substituted aryl; M is hydrogen, sodium, potassium, ammonium,

diethanolamine, cyclohexylamine, a naturally-occurring amino acid of MW less than 500 kD, lower alkyl (C₁₋₆), cycloalkyl, or aryl (C₆₋₁₂); and n is 0-5;

c) [or] Formula IV:

Formula IV




wherein R and R₁ are independently selected from the group consisting of H, lower alkoxy (C₁₋₆), or lower alkyl (C₁₋₆); R₂ is selected from the group consisting of aryl (C₆₋₁₂) and substituted aryl; M is hydrogen, sodium, potassium, ammonium, diethanolamine, cyclohexylamine, a naturally-occurring amino acid of MW less than 500 kD, lower alkyl (C₁₋₆), cycloalkyl, or aryl (C₆₋₁₂); and n is 0-5; or,

d) any combination thereof.

17. A pharmaceutical composition for the treatment or inhibition of hypercholesterolemia or hypertriglyceridemia, comprising:

92 a therapeutically-effective amount of a compound of either Formula I:

a) Formula I



92
Antil

Formula III

